NSK-A-TS AND NSK-B-TS

NSK-A-TS and NSK-B-TS Tuning Standards





Tandem Mass Spectrometer (MS/MS) Tuning Standards, NSK-A-TS and NSK-B-TS, have been developed to complement quality assurance and quality control (QA/QC) procedures in the laboratory. Use MS/MS Tuning Standards to:

- Ensure MS/MS instrument is operating at peak sensitivity for analysis of amino acids and acylcarnitines prior to analysis.
- Monitor instrument sensitivity from analysis of the first dried blood spot (DBS) to the last, whether samples are from one or several microtiter plates, during and between analysis runs.
- Quickly locate the source of sensitivity loss during an analytical run or between batch analyses.
- Compare performance of multiple instruments within a laboratory or across many laboratories.
- Evaluate performance before and after instrument maintenance.
- Assess MS/MS performance in analysis of amino acids (AA) and acylcarnitines (AC) independent of DBS samples and their preparation.

NSK-A-TS		
Catalog No.	Amino Acid	Conc. (µM)*
DLM-250	L-Alanine (D <sub>4</sub> )	25
CLM-1055	L-Phenylalanine (ring- <sup>13</sup> C <sub>6</sub> )	25
DLM-3860	L-Citrulline (5,5-D <sub>2</sub> )	25
DLM-335	DL-Glutamic acid (2,4,4-D <sub>3</sub> )	25
DLM-431	L-Methionine (methyl-D <sub>3</sub> )	25

	_	
	Amount	Price
NSK-AB-TS	One vial of each	\$480
(also available as individual vials)	Three vials of each	\$1,368
marriadar viais,	Five vials of each	\$2,160



After reconstitution in mobile phase whether as free acids or derivatized as butyl esters, the tuning standards are stable in solution for up to 30 days when stored at 4°C. The prepared solutions are ready for use immediately whether for tuning the instrument as part of regular maintenance, for troubleshooting MS/MS instrument problems or for a quick daily check before each batch run (or as often as a protocol may require). These reconstituted tuning standards are concentrated solutions and do not replace NSK-A and NSK-B reference standards.

NSK-B-TS		
Catalog No.	Carnitine	Conc. (µM)*
DLM-3555	L-Carnitine (D <sub>9</sub> , 98%) (CN)	7.6
DLM-3973	O-Propionyl-L-carnitine:HCl (D <sub>3</sub> ) (C3)	0.38
DLM-755	O-Octanoyl-L-carnitine:HCl (D <sub>3</sub> ) (C8)	0.38
DLM-1263	O-Palmitoyl-L-carnitine:HCl (D <sub>3</sub> ) (C16)	0.76

<sup>\*</sup>When reconstituted in 1 mL solvent.

For research use only. Not for diagnostic purposes.

To place an order please contact CIL:

t: 978.749.8000

1.800.322.1174 (North America)

cilsales@isotope.com